

# Amvisc® Plus

1.6% Sodium Hyaluronate

VISCOELASTIC



The viscoelastic that provides cohesive versatility at every stage of the cataract procedure.

## VERSATILE COHESION



## CRYSTAL-CLEAR VIEW

**A cohesive + a dispersive viscoelastic in one syringe to assist with every stage of the procedure.**

- Supports the bag fully for IOL delivery while allowing efficient haptic placement and lens rotation.<sup>1</sup>
- Cohesive properties ensure continuous space maintenance while providing corneal endothelial cell protection.<sup>1</sup>
- Cohesive versatility allows you to complete the procedure without needing a second viscoelastic.<sup>1</sup>

**Amvisc® Plus**  
1.6% Sodium Hyaluronate

**BAUSCH + LOMB**

# Amvisc® Plus


1.6% Sodium Hyaluronate

60081L order number 60081LXXXX

LEARN MORE AT [bauschsurgical.com](http://bauschsurgical.com)



MODEL NUMBER	SIZE	VISCOSITY, CP (AT 1S-1SHEAR RATE, 25°C)	MOLECULAR WEIGHT (DALTONS)	COMPOSITION	COHESION	OSMOLALITY (MOSM/KG)	CANNULA SIZE
60081L	0.8mL	55,700	>80,000	1.6%HA	Medium/Cohesive	340	27G



## PRODUCT NUMBER 60081L

### DESCRIPTION

- 1.6% sodium hyaluronate (HA) concentration
- 0.8-mL preloaded sterile glass syringe with finger grip for comfort
- 27-gauge disposable cannula, 32° angle
- Syringe requires assembly of the cannula only
- Not made with natural rubber latex
- >80,000 molecular weight (Daltons)

### DISPENSED

Singles

1. Amvisc® Plus Directions for Use

**STORZ**  
Ophthalmic Instruments

Find B+L IOL surgical equipment  
online at [www.StorzEye.com](http://www.StorzEye.com)

## Indications and Important Safety Information for Amvisc® and Amvisc® Plus Viscoelastics

**INDICATIONS:** Amvisc® and Amvisc® Plus are indicated for use as a surgical aid in ophthalmic anterior and posterior segment procedures including: Extraction of a cataract, implantation of an intraocular lens (IOL), corneal transplantation surgery, glaucoma filtering surgery, and surgical procedures to reattach the retina.

Because of its lubricating and viscoelastic properties, transparency, and ability to protect corneal endothelial cells, Amvisc and Amvisc Plus helps maintain anterior chamber depth and visibility, minimizes interaction between tissues, and acts as a tamponade and vitreous substitute during retinal reattachment surgery. Amvisc and Amvisc Plus also preserves tissue integrity and good visibility when used to fill the anterior and posterior segments of the eye following open sky procedures.

**PRECAUTIONS:** There may be increased intraocular pressure following surgery caused by pre-existing glaucoma or by the surgery itself. An excess quantity of Amvisc and Amvisc Plus should not be used. Amvisc and Amvisc Plus should be removed from the anterior chamber at the end of surgery to prevent or minimize post-operative intraocular pressure increases (spikes). If the postoperative pressure increases above expected values, correcting therapy should be administered. Reuse of cannula should be avoided. Store at 2-8° C. Protect from freezing.

**ADVERSE EVENTS:** Transient postoperative inflammatory reactions were reported in clinical trials and oral and topical steroid preparations were administered. The best index of the degree of phlogistic response is the postoperative clarity of the vitreous cavity. Transient postoperative increase in intraocular pressure has been observed following the use of sodium hyaluronate in anterior segment surgery. On rare occasions postoperative reactions, including inflammation, corneal edema, and corneal decompensation have been reported.

**ATTENTION:** Refer to the Directions for Use labeling for a complete listing of indications, warnings and precautions, clinical trial information, etc.

**CAUTION:** Federal (USA) law restricts this device to the sale by or on the order of a physician.